

Bacteriology of Pelvic Inflammatory Disease among Women Attending the Gynecology Clinic at Kampala International University Teaching Hospital, Uganda

Petrus Baruti¹, Emmanuel Nzabandora¹ and Ezera Agwu²

¹Department of Obstetrics and Gynecology of Kampala International University, Uganda.

²Department of Microbiology, Kampala International University, Uganda.

ABSTRACT

Pelvic inflammatory disease (PID) is major health problem in developed and developing country involving more young women. It is associated with high rate of female reproductive health morbidity; it can complicate with ectopic pregnancy, infertility and chronic pelvic pain. A poor response therapy increases the likelihood of these complications; this could be due to an increase in antimicrobial resistant pathogens. This was a cross-sectional study conducted among women who attended gynecology clinic at Kampala International University Teaching Hospital. Consecutive enrolment of 324 participants who consented to participate was done daily until a required sample size was realized from November 2019 to January 2020. Structured questionnaires were used to collect data on associated factors; endocervical swab was taken from patient clinically diagnosed with PID. Culturing for colony characteristics followed by Gram stain was used for provisional identity of pathogenic bacteria. Further identification was done by a set of biochemical tests. Antibacterial susceptibility pattern of isolated bacterial pathogens was determined by Kirby Bauer disc diffusion method, a rapid diagnostic test to detect Chlamydia antibody in the endocervical swab sample was also used to identify the Chlamydia trachomatis carriers among the patients. Data was analyzed using STATA VERSION 14.2. The isolated bacteria was *Staphylococcus aureus* and the least-prevalent pathogen was *Citrobacter species*. *Staphylococcus aureus* showed high susceptibility to Imipenem. The high resistance rate of more than 60% was seen to the commonly prescribed antibiotics as recommended by Uganda clinical guideline these were Doxycillin, ciprofloxacin, Cefixim and Ceftriaxon. *Staphylococcus aureus* is the commonest organism isolated from the endocervical swab of patients with PID. Awareness amongst health workers and patients about these major factors so that management can be directed. Rational use of antibiotics by health workers is paramount to combat resistance.

Keywords: Bacteriology, PID, Women, inflammation, Gynaecology

INTRODUCTION

Pelvic inflammatory disease is an infection which affects the upper female reproductive tract from the internal cervical opening, the uterus, fallopian tube, ovaries and pelvic peritoneum [1, 2, 3, 4, 5]. Its distribution and associated factors, management and control are challenges facing the gynecologists in an effort to discharge their corporate responsibilities to the general public [6, 7, 8, 9]. The oldest and clear description of pelvic inflammatory disease was done by Mauriceau in 1693 when he described puerperal infection with abscess in both

sides of the uterus [10, 11, 12, 13, 14]. While the first description of the bacterial agent that causes pelvic inflammatory disease was after Neisser discovered *Neisseria gonorrhea* in 1879 [15, 16]. This description was done by Westermarck in 1886 when he demonstrated the presence of *Neisseria gonorrhea* in tubal pus and Wertheim in 1894 the first to demonstrate these organisms invading the tube [17, 18, 19]. Chronic pelvic inflammatory tumors unconnected with the puerperal state were described by Simpson in 1843; which described the condition as pelvic cellulitis

[20]. *Neisseria Gonorrhea* was since then known as the leading cause of pelvic inflammatory disease (PID) and its history is strongly linked to the one of pelvic inflammatory disease. It was thought to be caused solely by an inflammation occurring in the cellular tissue of the pelvis [2].

RESEARCH METHODOLOGY

Study design

This was a cross sectional study. Laboratory investigations were done to achieve the prevalence, the common bacteria isolates and antibacterial susceptibility pattern in women with pelvic inflammatory disease attending gynecology clinic at Kampala international university teaching hospital. Association between PID and different factors was established.

Study area

The study was conducted at Kampala International University Teaching Hospital found in Ishaka Bushenyi Municipality at approximately 60km from Mbarara town along Mbarara Kasese highway.

Study population

The study populations were all women of reproductive age in the catchment area

Target population

All women of reproductive age attending gynecology clinic at Kampala international university teaching hospital shall be considered for inclusion in this study.

Inclusion criteria

All the women at the reproductive age attending gynecology clinic of Kampala international university teaching hospital as well as emancipated minors.

Exclusion criteria

Women on antibiotics, pregnant women, unconscious patients who cannot consent and minors were excluded from the study.

Sample size determination

Objective number 1, the sample size was determined using [3] formula with the

Purpose of the study

To determine the common bacterial pathogens and antibacterial susceptibility pattern among women with pelvic inflammatory disease attending gynecology clinic at Kampala University International Teaching Hospital.

estimated prevalence of 50%, because of lack of current prevalence of PID in Uganda.

$$n = \frac{z^2 pq}{d^2}$$

Where;

n = Desired sample size

z = Standard normal deviation at 95% level of confidence;

z= 1.96

p = Prevalence of pelvic inflammatory disease in Uganda, assumed at 50%, and

d= Level of precision=0.05

$$n = \frac{(Z_{\alpha})^2 \times p(1 - p)}{(d)^2}$$

$$n = \frac{(1.96)^2 \times 0.5(1 - 0.5)}{(0.05)^2}$$

N₁= 384 this is considered as an assumption of sample size.

Objective number 2:

Musa and colleagues [4] in the study done in Mbarara Regional Referral Hospital on screening for *Chlamydia Trachomatis* among women of reproductive, found that 26% of women were carriers of *Chlamydia Trachomatis*.

$$n = \frac{z^2 pq}{d^2} \quad n = \frac{(Z_\alpha)^2 \times p(1-p)}{(d)^2}$$

Where;

n = Desired sample size

z = Standard normal deviate at 95% level of confidence; z= 1.96

p = prevalence of Chlamydia trachomatis in reproductive aged women, using a study done in Mbarara by Musa and colleagues [4]

d=Level of precision= 0.05

$$n = \frac{(1.96)^2 \cdot 0.2(1-0.2)}{(0.05)^2}$$

N₂= 295

Specific objective 3: Mark *et al.* [5] reported that women with a previous history of PID have a greatest risk of having subsequent PID, OR = 5.9, the proportion (p) PID among woman with previous history of PID 13.1% and among women without previous history of PID was 3.6%

r=ratio of previous history of PID to those without previous history of PID, r=13.1:3.6, r=3.6:1=3.6

Where; =z-statistic at α=0.05, = z-statistic at β=0.84 hence statistical power = 80%.

Using the

$$n = \frac{(1+r)^2 [Z_\alpha + Z_\beta]^2}{r (\ln OR)^2 \times p(1-p)}$$

Formula [6]: where

$$\frac{(1+3.6)^2 \times (1.96+0.84)^2}{3.6(\ln 5.9)^2 \times 0.131(1-0.131)}$$

n₃=118

Decision on the sample size:

Sample size for objective number 2 was considered since this was a research done in Uganda (Mbarara Regional Referral hospital) and reflects the true Ugandan reality. 10% of this sample size was added to minimize non response or loss of some laboratory request form, therefore the sample size was:

N₂=324

Sampling technique

Consecutive sampling method was used to select participants who consented to be part of the study. All the women of reproductive age who met the inclusion criteria was invited to participate in the study, the participants was enrolled according to their order of arrival in gynecology clinic and this was carried out on a daily basis until the required sample size was achieved.

Data collection instruments

A pretested questionnaire was administered to each participant who consented to participate to the study in order to collect information on socio-demographic, gynecological and sexual behavior factors that related to the development of pelvic inflammatory disease in. A detailed history was taken in English, translated in local language where necessary for women who could not understand English physical examination was carried out and the endocervical sample was taken from patient with symptoms and of PID in order to achieve all the objectives.

Sample processing and analysis

Isolation

Samples collected using a sterile procedure with the endocervical swab stick was inoculated on blood agar, chocolate agar, Mac Conkey agar, Thayer Martin medium, and different biochemical tests were used. After, they were incubated both aerobically and anaerobically at 37°C for 24-48hrs. Colony morphology were observed according to shape, size, elevation, margin and surface characteristics.

Rapid diagnostic test was used in order to identify the *Chlamydia trachomatis* antibody careers within the endocervical sample of the participants, the isolation of Chlamydia which uses living cells (McCoy cell) was not done due lack of this specific

media to culture *Chlamydia trachomatis*, this rapid Chlamydia test was used to determine the percentage of Chlamydia carriers among the patient with pelvic inflammatory disease.

Direct gram microscopy

A direct smear was made for Gram stain according to [7]; a drop of sterile normal saline was added on the center of a clean dried glass slide and the swab containing the sample rolled in the drop of normal saline spreading it on the glass slide in a circular motion to make a thin smear. The smear was then allowed to air dry and then heat fixed by passing it at least three times over a Bunsen flame. The slide was placed on the staining rack and flood with crystal violet solution for 60 seconds, wash with clean water and cover with lugol's Iodine and then it was allowed to act for a minute. The slide was washed in clean water and then decolorized with 50% acetone alcohol under slow running tap water until a faint pink color is observed or no more color tend to flow from the smear. The process of discoloration was not exceeding 30 seconds.

After decolorizing, it was washed in clean water and counter stain with neutral red solution. The slide was again washed in clean water; air dried and observed under the microscope with X100 objective lens (oil immersion lens).

Gram positive bacteria were observed as blue or purple color and Gram negative as red or pink color. Also, the morphology and shape of the bacteria were identified whether they are cocci, diplococcic, cocci in chains, clusters, and whether they are rods in appearance [7].

Identification of bacterial isolates

Biochemical tests

The isolates were identified using the conventional biochemical tests such as catalase, optochin, coagulase, indole, citrate utilization, urea utilization, triple sugar iron agar fermentation, MR-VP test and oxidase as described below;

Catalase test

Catalase test was carried out according to the method described by [7], to determine the ability of the isolate to produce the enzyme, catalase. A drop of 3% hydrogen peroxide was added to a loop full of the

test organisms. Presence of bubbles indicated catalase activity.

Indole test

Indole test was carried out according to the method described by [7] to determine the ability of the isolate to degrade amino acid tryptophan and produce tryptophanase, enzyme were tested. 1% tryptophan broth in a test tube were inoculated with isolate and incubated at 37°C for 48 hours. After 48 hours, 1 milliliter of chloroform was added to the broth. The test tube was shaken gently, and 2.1 ml of Kovac's reagent were added and again shaken gently, this was allowed to stand for 20 minutes. The formation of red coloration at the top layer, indicated positive while yellow coloration indicated negative.

Urease test

Urease test was done according to the method described by [7] to determine the ability to hydrolyze urea to produce ammonia and carbon dioxide. Test organism was inoculated into urease broth and incubated at 30°C for 72 hours. Purplish pink coloration of the medium indicates positive reaction.

Citrate utilization

This was carried out by inoculating the test organism in test tube containing Simon's citrate medium and incubated for 24 to 72 hours. The development of deep blue color after incubation indicated a positive result [7].

Triple sugar iron test

Triple sugar iron test was carried out according to the method described by [7]; the test determines the ability of the organism to ferment the three sugar component of the medium: glucose, lactose and sucrose. The medium contains a pH indicator (phenol red) and a detection system (thiosulphate and ferrous sulphate) for hydrogen sulphide (H₂S).

The medium was prepared as an agar slant. The test organism was inoculated by stabbing the medium using sterilized straight wire loop and the surface of the slope were also streaked with the test organism. The tests were incubated at 37°C for 72 hours. After incubation, gas production was determined by observing the cracking of the medium, and

production of H₂S was observed by the blackening of the bottom of the medium.

Coagulase test

This test was used to identify *Staphylococcus Aureus* which produces the enzyme coagulase. The rapid slide test was done by placing a drop of distilled water on each end of slide. Then a colony of the test organism (previously checked by Gram staining) was emulsified in each of the drops to make two thick suspensions.

A loopful of plasma was added to one of the suspensions and mixed gently. Formation of clumps of the organisms within 10 seconds was suggestive of a positive test while absence of these clumps indicated negative results. For suspected *Staphylococcus aureus* isolates which turn negative for the rapid slide test, tube test was done by emulsifying several isolated colonies of test organism in 1 ml of diluted rabbit plasma (1:5) dilution to give a milky suspension. The tubes were then incubated at 35°C in water bath for 4 hours. These were examined at intervals of 1, 2 and 4 hours for clot formation by tilting the tube through 90°. If the test is still negative, the tube was left at room temperature overnight and examined again for *Staphylococcus aureus* that produce a delayed clot [7].

Oxidase test

The test was used in identification of organisms which produce the enzyme cytochrome oxidase. A filter paper soaked with the substrate tetra methyl-p-phenylenediamine Dihydrochloride was moistened with sterile distilled water. Using a piece of stick or glass rod, a colony of the test organism was smeared on the filter paper. The development of a blue-purple color within a 10 seconds indicated positive test while absence or formation of a blue-purple color after 10 seconds was considered negative [7].

Susceptibility pattern determination (Kirby-Bauer disc diffusion technique)

Pure culture colonies of the organisms was tested for their antimicrobial drug susceptibility against twelve commonly used antibiotics including ceftriaxone (30ug), Ampicillin (30ug), Cefixim(30ug), Ofloxacin (30ug), Levofloxacin(30ug), gentamycin, Ciprofloxacin, nitrofurantoin,

erythromycin, among others using Kirby-Bauer disc diffusion (Banuer AW, 1985) as modified by the clinical laboratory standard institute (CLSI, 2010). Selection of 3 to 5 isolated colonies from the medium were done using a sterile wire loop and transferred into 3mls of nutrient broth in a bijou bottle. The broth was mixed by inversion for complete dissolution of the colonies. The mixture was incubated for 4 minutes and was compared with 0.5% McFarland turbidity standard. A sterile inoculation glass rod was used to spread the surface of Muller-Hinton agar plates homogenously with the diluted colonies. Antibiotic discs were placed onto the inoculated Muller-Hinton agar and incubate at 37°C for 24-48 hours. The plates were examined and the diameter of zones of inhibition was measured in mm using a meter ruler and was compared to a standard chart of the corresponding antibiotics used for measuring zone of inhibition. Zone of inhibition measured recorded as susceptible (S), Intermediate (I), and Resistant (R) according to the standard chart [7].

Data analysis plan

Data from questionnaires were entered in Microsoft Excel 2010, and thereafter exported to STATA 14.1. Socio-demographic, sexual behaviors and gynecologic factors were summarized as means and medians, standard deviations and interquartile range (for continuous variables) were determined. Proportions, percentages and frequencies were used for categorical variables using STATA 14.1.

Ethical considerations

Informed consent

Informed consent and respect for participant's voluntary recruitment was observed. Informed consent for participants were obtained and signed after fully explaining the details of the study to them in English and local languages where necessary (copy attached at Appendix). Participants were not forced to enroll themselves if they don't want to, they were free to withdraw from the study any time they wish without coercion or compromise of care they are entitled to.

Risks and adverse events to study participants

Patients may undergo pain during swabbing and speculum examination, however, the process of obtaining a swab was done gently and professionally to minimize risk of pain and minimize re-infection as far as possible. Additionally, culture and sensitivity are the recommended guidelines prior to antibiotic therapy to minimize the risk of antibiotic resistance.

Approval procedure

Approval to carry out the study shall be sought from the department of obstetrics and gynecology, the faculty and post graduate directorate and finally, Research Ethics Committee of Kampala International University. This approval letter shall be presented to the hospital administration of KIU-TH. Permission shall be sought from the administration of the hospital before the study is conducted. The study will be registered with Uganda National Council for Science and Technology.

RESULTS**Socio demographic findings****Table 1: Socio demographic factors**

Characteristics	Frequency	%
Age (years)		
<20	31	9.6
20-29	205	63.3
30-39	71	22.0
40-49	17	5.1
Education		
None	11	3.4
Primary	99	30.6
Secondary	111	34.4
Tertiary	103	31.6
Occupation		
None	127	39.2
Farmer	85	26.2
Professionals	51	15.7
Business	31	9.6
Manual laborer	30	2.3
Monthly income (UGX)		
None	10	3.1
<300000	230	71.8
300000-600000	66	26.5
>600.000	18	5.6
Marital status		
Single	86	26.5
Married	238	73.5

The above table illustrates that 63.3% of participants are aged of 20-29 years, 34.4% have secondary education, 39.2% have no

occupation, 71.8% of participants have a monthly income of less than 300.000 Uganda Shillings and 73.5% are married.

Table 2 Gynecological factors

Characteristics	Frequency	%
Parity		
Zero	98	30.3
1-3	153	47.2
>3	73	22.5
Had PID before		
No	224	69.1
yes	100	30.9
Had miscarriage before		
No	264	81.5
Yes	60	18.5
Use Contraceptive		
No	132	40.7
yes	192	59.3
Intra Uterine Procedure		
No	281	86.7
Yes	43	13.3
Type contraception		
Condoms	38	19.8
Pills	61	31.8
Injectables	65	33.8
IUD	28	14.6
Type of miscarriage		
Spontaneous	41	68.3
induced	19	31.7

From the above table, 47.2% of the study participants had delivered at least one to three times, 69.1% had had miscarriage of which 68.3% were spontaneous, 59.3% of the study participants had ever used

contraceptive methods of which 33.6% had used injectable contraceptive methods and 86.7% had not had intrauterine procedures.

Table 3: Sexual behavior factors

Characteristics	Frequency	Percent
Number of of sexual partners		
None	20	6.2
One	253	78.0
More than one	51	15.8
Age of initiation sexual activity(year)		
< 15	25	7.7
16-20	242	74.7
>20	57	17.6
Condom Use		
Sometimes	84	25.9
Every time	38	11.8
Never	202	62.3
Smoking		
Never smoke	316	97.5
Ever smoke	8	2.5

The above table shows that, the age of initiation of sexual activity for the majority

of participants was 16-20years in 74.7%, most of the study participants denied the

use of condoms with 62.3% and 97.5% were nonsmokers.


Table 4: Prevalence of the isolated bacteria

Bacteria I Isolates	Fr(%)
Staphylococcus Aureus	24(38.7)
Chlamydia trachomatis	19(30.7)
Neisseria species	12(19.7)
Streptococcus species	10(16.4)
Pseudomonas species	03(4.8)
Escherichia coli	03(4.8)
Klebsiella	32(3.2)
Gardenella species	01(1.6)
Citrobacter species	01(1.6)

From the above table, *Staphylococcus aureus* was the most prevalent bacteria isolated with 38.7% followed by *Chlamydia trachomatis* which was indirectly identified using Chlamydia rapid

diagnostic test with 30.7%, *Neisseria* and *streptococcus* were isolated in the proportion of 19.6% and 16.4% respectively.

Figure 1: Isolated microorganisms



From the above a total eight microorganism were identified, the most common being *staphylococcus Aureus* with 38.7% followed by Chlamydia identified

indirectly with 30.7%, *Neisseria* and *Streptococcus* were in respectively 19.4% and 16.1%.

Table 5: Susceptibility of Common bacteria isolates

Drug	Bacteria Isolates		
	Staph Aureus	Neisseria spp	Streptococcus spp
Doxycilin	00	02(16.7)	02(20)
Cefixim	00	00	00
Ampicillin	00	00	02(20)
Ciprofloxacin	1(4.4)	00	1(10)
Azithromycin	2(8.3)	05(41.7)	05(50)
Levofloxacin	3(12)	2(16.7)	1(10)
Ceftriaxone	3(12)	2(16.7)	00
Nitrofurantoin	12(50)	1(8.3)	00
Vancomycin	13(54)	00	5(50)
Imipenem	19(79)	08(66.)	9(90)
Gentamycin	12(50)	06(54.6)	4(44.4)

The above table shows the antibacterial susceptibility pattern against the common bacterial isolates among women with pelvic inflammatory disease. The most common bacteria are sensitive to

Imipenem at 79.1% for *staphylococcus Aureus*, *Neisseria species* and *streptococcus species* were also sensitive to imipenem with respectively 66.7 and 90%.

Table 6: Susceptibility and Resistance pattern of the common bacteria isolates among women with PID at KIU-TH


	Staphylococcus aureus			Neisseria Species			Streptococcus species		
Drugs	S	I	R	S	I	R	S	I	R
Doxycilin	00	7(29.2)	17(70.8)	02(16.7)	00	10(83.3)	02(20)	00	8(80)
Cefixim	00	6(25)	18(75)	00	00	10(100)	00	00	10(100)
Ampicill	00	4(17.4)	19(82.6)	00	(2)16.7	10(83.3)	02(20)	00	80%
Ciprofloxacin	1(4.4)	5(21.7)	17(73.9)	00	2(16.7))	10(83.3)	1(10)	00	9(90)
Azithromycin	2(8.3)	13(54.2)	9(37.5)	05(41.7)	6(50)	1(8.3)	05(50)	5(50)	00
Levofloxacin	3(12.5)	4(16.7)	17(70.8)	2(16.7)	2(18.2)	5(45)	1(10)	00	9(90)
Ceftriaxone	3(12.5)	9(37.5)	12(50)	2(16.6)	1(8.33)	9(75)	00	3(30)	00
Nitrofurantoin	12(50)	8(33.3)	4(16.7)	1(8.3)	4(33.3)	7(58.3)	5(50)	3(30)	2(20)
Vancomycin	13(54)	8(33)	3(12.5)	00	9(75.0)	3(25)	5(50)	4(40)	1(10)
Imipenem	19(79)	00	5(21)	08(66.7)	00	4(33.3)	9(90)	1(10)	00
Gentamycin	12(54.6)	3(13.6)	7(31.8)	6(54.6)	1(9.1)	4(36.4)	4(44.4)	00	5(55.6)
	Staphylococcus aureus			Neisseria Species			Streptococcus species		
Drugs	S	I	R	S	I	R	S	I	R
Doxycilin	00	7(29.2)	17(70.8)	02(16.7)	00	10(83.3)	02(20)	00	8(80)
Cefixim	00	6(25)	18(75)	00	00	10(100)	00	00	10(100)
Ampicill	00	4(17.4)	19(82.6)	00	(2)16.7	10(83.3)	02(20)	00	80%
Ciprofloxacin	1(4.4)	5(21.7)	17(73.9)	00	2(16.7))	10(83.3)	1(10)	00	9(90)
Azithromycin	2(8.3)	13(54.2)	9(37.5)	05(41.7)	6(50)	1(8.3)	05(50)	5(50)	00
Levofloxacin	3(12.5)	4(16.7)	17(70.8)	2(16.7)	2(18.2)	5(45)	1(10)	00	9(90)
Ceftriaxone	3(12.5)	9(37.5)	12(50)	2(16.6)	1(8.33)	9(75)	00	3(30)	00
Nitrofurantoin	12(50)	8(33.3)	4(16.7)	1(8.3)	4(33.3)	7(58.3)	5(50)	3(30)	2(20)
Vancomycin	13(54)	8(33)	3(12.5)	00	9(75.0)	3(25)	5(50)	4(40)	1(10)

Imipenem	19(79)	00	5(21)	08(66.7)	00	4(33.3)	9(90)	1(10)	00
Gentamycin	12(54.6)	3(13.6)	7(31.8)	6(54.6)	1(9.1)	4(36.4)	4(44.4)	00	5(55.6)

From the above table, the common bacteria isolates among women with PID exhibit a high resistance rate of more than 70%

towards the commonly used antibacterial like Doxycillin, cefixim, ampicillin and ciprofloxacin.


Figure 2: Antibacterial susceptibility of Staphylococcus Aureus



From the graph above, *Staphylococcus Aureus* is most sensitive to Imipenem with 79.2%, followed by Gentamycin,

Vancomycin and Nitrofurantoin with respectively 54%, 54% and 50%.


Figure 3: Antibacterial susceptibility to Neisseria species



The Above graph shows that *Neisseria species* are most sensitive to Imipenem with 96%, followed by ceftriaxon with 56%,

gentamycin was sensitive in 54%, Azithromycin and Ofloxacin represented respectively 41% and 40%.

Figure 4: Antibacterial susceptibility to Streptococcus species



This above graph illustrates that *Streptococcus* species were more sensitive to Imipenem at 90%, followed by

According to this study *Staphylococcus aureus* was the most common isolated bacterium accounting for 37.8% of PID cases, followed by *Chlamydia trachomatis* and *neisseria species* with respectively 30% and 16%. The study of [8] in Cameroon found that common bacteria isolates were Genital tract *mycoplasma* with 54.3% followed by *Chlamydia trachomatis* with 37.1% this was high compare our study, he also reported a low prevalence of *Staphylococcus aureus* with 2.9%.

This result was also higher to Spencer who found a prevalence of 16% for *Staphylococcus aureus* which was followed by *E. coli* 12% and *Streptococcus species* 8%. The prevalence of *Chlamydia trachomatis* in this was 30%, this was higher to the one found by Musa and colleague in Mbarara regional referral Hospital (2017) which was 26.4%.

nitrofurantoin with 50%, Azithromycin with 50%, Gentamycin and Ceftriaxone had respectively 44.5 and 30%.

DISCUSSION

In this study, the most common bacterial isolate were *staphylococcus Aureus*, *Neisseria species* they were all susceptible to Imipenem at respectively 79.8%, 66.7% and 90%, this result differ from Obluiru in Nigeria *Staphylococcus* was more sensitive to ceftazidin, It also disagree with Spencer and colleagues (2014) where cefotaxim was established as the sensitive drug for the treatment of PID. *Staphylococcus aureus* exhibited susceptibility to nitrofurantoin and gentamycin in 50% of cases each in this study, this has been similarly observed by Spencer and colleagues where gentamycin showed the same inhibitory affect against *Staphylococcus aureus*.

Neisseria species were more susceptible to Imipenem at 66.7% followed by gentamycin and Azythromycin with respectively 54.6% and 41.7% this disagree with [9] in Cameroon who established ceftriaxon as a drug of choice in PID

against which *Neisseria* had a resistance of 1% as showed in their study. The susceptibility of *Chlamydia* was not established in this study.

Streptococcus was found to more sensitive to Imipenem in 90% of cases, it also showed a susceptibility to gentamycin and Azithromycin of 50% this was contrary to

the findings of [10] were streptococcus species were more sensitive to Erythromycin and Azithromycin [10]. This study demonstrates the high resistance rate to the commonly used antibiotic in case of PID in outpatient clinic at KIU-TH. These are ceftriaxon, cefixim, levofloxacin and Doxycillin.

CONCLUSION

The common bacteria pathogens isolated were *Staphylococcus Aureus*, *Chlamydia Trachomatis*, *Neisseria species* and

Streptococcus species; Imipenem was the most sensitive antibiotic for both of the bacteria.

REFERENCES

1. Viberga, I. (2006). *The Clinical Appearance of Pelvic Inflammatory Disease in Relation to Use of Intrauterine Device in Latvia*.
2. Rees, E., Annels, E. H. and Both, B. (1969). Gonococcal salpingitis*.
3. Susan P. Hamill, (2015). *Untangling the Mystery of Teaching Business Organizations*, 59 ST. LOUIS U. L.J. (2015).
4. Musa, M., Joel, B., Lenard, A., Joseph, N., Ronald, M. and Julius, M. (2016). Prevalence and Factors Associated With Genital Chlamydial Infections among Women Attending the Gynaecology Clinic At Mbarara Regional Referral Prevalence and Factors Associated With Genital Chlamydial Infections among Women Attending the Gynaecology Clinic At Mbarara Regional Referral Hospital, (May).
5. Marks, C., Tideman, R. L., Estcourt, C. S., Berry, G. and Mindel, A. (2000a). Assessment of risk for pelvic inflammatory disease in an urban sexual health population, 470-473.
6. Suresh, K. P. and Chandrashekara, S. (2012) Sample Size Estimation and Power Analysis for Clinical Research Studies. *Journal of Human Reproductive Science*, 5, 7-13.
7. Cheesbrough, M. (2006) *District Laboratory Practice in Tropical Countries*. Part 2, 2nd Edition, Cambridge University Press Publication, South Africa, 1-434.
8. Nkwabong, E. and Dingom, M. A. N. (2015). Acute Pelvic Inflammatory Disease in Cameroon: A Cross Sectional Descriptive Study, 19(December), 87-91.
9. Tayimetha, C. Y., & Unemo, M. (2018). Antimicrobial Susceptibility of *Neisseria gonorrhoeae* Isolates in Yaoundé , Cameroon From 2009 to 2014, 45(12), 101-103. <https://doi.org/10.1097/OLQ.0000000000000915>
10. Ssempijja Fred, Marta Vicente-Crespo, Dare Samuel Sunday and Edmund Eriya Bukenya (2022). The effect of *Imperata cylindrica* (L.) P. Beauv. on Brain tissue Morphology of *Drosophila Melanogaster transgenic* flies overexpressing the paralytic Gene. *IDOSR Journal of Biochemistry, Biotechnology and Allied Fields* 7(1): 1-11.
11. Ssempijja Fred, Marta Vicente-Crespo, Dare Samuel Sunday1 and Edmund Eriya Bukenya (2022). The Effect of *Imperata Cylindrica* (L.) P. Beauv on Learning and memory performance in a single and prolonged exposure treatments of *Drosophila Melanogaster transgenic* flies overexpressing the paralytic gene. *IDOSR Journal of Biochemistry, Biotechnology and Allied Fields* 7(1): 12-22
12. Ssempijja Fred, Marta Vicente-Crespo, Dare Samuel Sunday and Edmund Eriya Bukenya (2022). The Effect of *Imperata cylindrica* (L.) P. Beauv. on Myelin and Nissl Substance of *Drosophila melanogaster transgenic* Flies Overexpressing the Paralytic Gene. *IDOSR Journal of Biology, Chemistry And Pharmacy* 7(1)1-13.
13. Hussein Osman Ahmed, Joy Muhumuza and Musiime James Nabaasa

- (2022). Evaluation of the factors associated with immediate adverse maternal outcomes among referred women in labor at Kampala International University Teaching Hospital. IAA Journal of Biological Sciences 8(1):228-238
14. Hussein Osman Ahmed, Joy Muhumuza and Musiime James Nabaasa (2022). Factors associated with Immediate Adverse Maternal Outcomes among Referred Women in Labor attending Kampala International University Teaching Hospital. IAA Journal of Applied Sciences 8(1):117-125.
 15. Hussein Osman Ahmed, Joy Muhumuza and Musiime James Nabaasa (2022). The composite immediate adverse maternal outcomes among women in labor referred to Kampala International University Teaching Hospital IAA Journal of Scientific Research 8(1):149-156.
 16. Daniel Asimwe, Herman Lule and Izimba Daniel (2022). Epidemiology of Assault Injuries among Trauma Patients Presenting at Kampala International University Teaching Hospital and Jinja Regional Referral Hospital. INOSR Applied Sciences 8(1):111-119.
 17. E.O.Ikuomola,O.S Akinsonmisoye, R.O. Owolabi and M. B. Okon (2022). Assessment of Toxicity Potential of Secnidazole on Reproductive System of Male Wistar Rats. *INOSR Applied Sciences* 8(1):120-133
 18. Ugwu Okechukwu, P. C., Onwe, S. C. and Okon, M. B. (2022). The effect of Methanol Extract of *Rauwolfia vomitoria* on Lipid Profile of Chloroform intoxicated Wistar Albino Rats. IAA Journal of Scientific Research, 8 (1), 73-82
 19. E.O.Ikuomola,O.S.Akinsonmisoye, R.O. Owolabi and M. B. Okon (2022).Evaluation of the Effect of Secnidazole on Sperm Motility, Morphology, Viability and Total Sperm Count of Wistar Rats. *INOSR Experimental Sciences* 8(1): 74-83, 2022.
 20. E.O. Ikuomola , O.S Akinsonmisoye, R.O. Owolabi and M. B. Okon (2022).Evaluation of the effect of secnidazole on the histology of the testes and epididymis of male Wistar rats. *INOSR Experimental Sciences* 8(1): 84-94, 2022.